

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 215/28, 213/08, 22/04	A1	(11) International Publication Number: WO 00/27798 (43) International Publication Date: 18 May 2000 (18.05.00)
(21) International Application Number: PCT/EP99/08577 (22) International Filing Date: 9 November 1999 (09.11.99) (30) Priority Data: 9824706.7 11 November 1998 (11.11.98) GB 9824705.9 11 November 1998 (11.11.98) GB (71) Applicant (for all designated States except AT/US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (71) Applicants (for all designated States except US): TAITO CO. LTD. [JP/JP]; 7-5, Nihonbashi Odenma-cho, Chuo-ku, Tokyo 103-0011 (JP). YOSHITOMI PHARMACEUTICAL INDUSTRIES LTD. [JP/JP]; 6-9, Hiranomachi 2-chome, Chuo-ku, Osaka 541-0046 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): ABEL, Stephan [DE/DE]; Alemannenweg 12, D-79576 Weil (DE). FUJITA, Tet-suro [JP/JP]; 40-23, Daigokuden, Kaide-cho, Muko-shi,		Kyoto 617-0004 (JP). HIROSE, Ryoji [JP/JP]; Royal-court Nishi-akashi 2-302, Toba 1399-1, Akashi, Hyogo 673-0003 (JP). JORDINE, Guido [DE/DE]; Lehener Strasse 148, D-79106 Freiburg (DE). MISHINA, Tadashi [JP/JP]; Sharrman Coop Kuzzha 1-102, Minami-Kuzuha 2-10-2, Hirakata-shi, Osaka 573-1105 (JP). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PRODUCTION OF 2-AMINO-2-[2-(4-C ₂₋₂₀ -ALKYL-PHENYL)ETHYL]PROPANE-1,3-DIOLS (57) Abstract Disclosed is a process for the production of 2-amino-2-[2-(4-C ₂₋₂₀ -alkyl-phenyl)ethyl]propane-1,3-diols.		


FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Production of 2-amino-2-[2-(4-C₂₋₂₀-alkyl-phenyl)ethyl]propane-1,3-diols

The present invention relates to a process for the production of 2-amino-2-[2-(4-C₂₋₂₀-alkyl-phenyl)ethyl]propane-1,3-diols, particularly 2-amino-2-[2-(4-octylphenyl)ethyl]propane-1,3-diol of formula




hereinafter defined as Compound A.

2-Amino-2-[2-(4-C₂₋₂₀-alkyl-phenyl)ethyl]propane-1,3-diols are disclosed in EP-A1-627,406 the relevant disclosure of which is incorporated herein by reference. On the basis of observed activity, e.g. as described in EP-A1-627,406, they have been found to be useful e.g. as immunosuppressants, e.g. in the prevention or treatment of acute allograft rejection or autoimmune diseases, or e.g. as described in WO 98/22100, in the prevention or treatment of xenograft rejection.

In accordance with the present invention, new processes for an improved production route of 2-amino-2-[2-(4-C₂₋₂₀-alkyl-phenyl)ethyl]propane-1,3-diols, particularly Compound A, in free form or in salt form, have now been found. It is to be understood hereinafter that the process steps of the invention may lead to 2-amino-2-[2-(4-C₂₋₂₀-alkyl-phenyl)ethyl]propane-1,3-diols in free form which may be converted into a salt form, or vice versa.


Accordingly, there is provided a process for preparing a 2-amino-2-[2-(4-C₂₋₂₀alkyl phenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises
g) reducing a compound of formula (20)

- 2 -



(20)


wherein R is C_{2-20} alkyl, each of R_1 and R_2 , independently, is C_{1-4} alkyl and R_3 is a protecting group,
 followed by a treatment with $(R_4-CO)_2O$ wherein R_4 is C_{1-4} alkyl, to obtain a compound of formula (21)



(21)

wherein R , R_3 and R_4 are as defined above; and

- i) treating the resulting compound of formula (21) with a basic compound to obtain a compound of formula (22)



(22)

wherein R is as defined above; and

- j) hydrogenating the resulting compound (22)

Any alkyl group mentioned herein may be a straight or branched chain alkyl group.
 Preferably C_{2-20} alkyl is octyl, particularly straight chain.

- 3 -


Examples of protecting groups as R_3 are e.g. disclosed in "Protective Groups in Organic Synthesis", T.W. Greene, J. Wiley & Sons NY (1981), 219-287, e.g. acyl such as formyl, acetyl, benzoyl; alkoxycarbonyl e.g. t-butyloxycarbonyl; allyloxycarbonyl; trityl; etc.

The reduction step g) may conveniently be performed using a reducing agent, e.g. NaBH_4 . The following treatment with an anhydride may be effected e.g. with acetic anhydride, preferably in the presence of a neutralizing agent, e.g. an aliphatic or organic amine, e.g. pyridine.

Suitable basic compounds for use in step i) include an alkali metal hydroxide, e.g. NaOH , KOH or LiOH . Step i) may preferably be carried out in a polar solvent, e.g. an alcohol such as methanol, at reflux. Step j) may be effected in accordance with known hydrogenation methods, e.g. in the presence of a hydrogenation catalyst, e.g. Pd-C .

As an alternative to above process, a 2-amino-2-[2-(4- C_{2-20} alkylphenyl)ethyl]propane-1,3-diol in free form or in salt form may be prepared by

- h) hydrogenating a compound of formula (21) as defined above to obtain a compound of formula (9)



wherein R , R_3 and R_4 are as defined above; and

- N) treating the resulting compound of formula (9) with a basic compound.

Steps h) and N) may be effected in accordance with methods known in the art, e.g. as disclosed above for steps j) and i), respectively, or in Examples 4 and 5 hereinafter.

According to a further embodiment of the invention, there is provided a process for preparing 2-amino-2-[2-(4- C_{2-20} alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises


- 4 -

- x) reducing a compound of formula (20) as defined above, followed by a treatment with $(R_4CO)_2O$ to obtain a compound of formula (9) as defined above; and
- N) treating the resulting compound of formula (9) with a basic compound.


In process step x) the reduction may comprise either a hydrogenation step, e.g. as disclosed above for step (j), followed by a reduction step, e.g. as disclosed above for step (g), or a reduction step as disclosed for step (g) comprising an intermediary hydrogenation step [e.g. as disclosed for step (j)] prior to the anhydride $(R_4CO)_2O$ treatment. Step (N) may be carried out as indicated above.

In a further or alternative embodiment, the invention provides a process for preparing 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises

- f) hydrogenating a compound of formula (20) as defined above or a compound of formula (20')



wherein R_1 , R_2 and R_3 are as defined above, and R' is C₁₋₁₉-CO- to obtain a compound of formula (8)



wherein R and R_1 to R_3 are as defined above; and

- K) reducing the resulting compound of formula (8) followed by a treatment with $(R_4CO)_2O$ to obtain a compound of formula (9) as defined above; and
- N) treating the resulting compound of formula (9) with a basic compound.


- 5 -

Process steps (f) and (K) may be performed as disclosed above for (j) and (g), respectively. Alternatively, the reduction step (K) may be carried out in the presence of $\text{Ca}(\text{BH}_4)_2$. The latter may be generated by reaction of NaBH_4 with CaCl_2 .

Compound of formula (20') may be prepared as disclosed in PCT/JP98/02998.


In a further or alternative embodiment, the invention further provides a process for preparing a 2-amino-2-[2-(4- C_{2-20} alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises

- g') reducing a compound of formula (20) as defined above under conditions to obtain a compound of formula (21')



wherein R and R_3 are as defined above; and

- y) hydrogenating the resulting compound of formula (21') to obtain a compound of formula (21'')



wherein R and R_3 are as defined above; and

- z) treating the resulting compound of formula (21'') with a basic compound.

Process step (g') is a reduction carried out in the same manner as in step (g). Suitable reducing agents may be e.g. as disclosed above for step (g). Steps (y) and (z) may be carried out as disclosed above for steps (j) and (N).

As an alternative to above step (y),

- i') a compound of formula (21') as defined above may be reacted with a basic compound, e.g. an alkali metal hydroxide, to obtain a compound of formula (22) as defined above, which compound of formula (22) is then further treated as disclosed above for step (j).


Suitable alkali metal hydroxide may be e.g. LiOH.

Compound of formula (20), used as starting product in the preparation of a 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, is novel and also forms part of the present invention, as well as its preparation. Compound of formula (20) may be obtained by a process comprising

- a) reacting a compound of formula (16)




wherein R is as defined above, with 2-bromoacetyl chloride to obtain a compound of formula (18)



wherein R is as defined above; and


- d) reacting the resulting compound of formula (18) with a compound of formula (2)



wherein R₁ to R₃ are as defined above.

- 7 -


Process step (a) may be performed in accordance with known bromoacetylation methods, e.g. in the presence of AlCl_3 . Process step (d) may conveniently be effected in the presence of sodium ethylate, in an anhydrous solvent such as anhydrous ethanol, and under an inert atmosphere, e.g. nitrogen, e.g. as disclosed in Example 13. Compound of formula (18) used in above step (d) may also be prepared by reacting a compound of formula (16) as defined above, with acetyl chloride to obtain a compound of formula (18')



wherein R is as defined above, which compound of formula (18') is then brominated.

In a further or alternative embodiment, the invention further provides a process for preparing a 2-amino-2-[2-(4- C_{2-20} alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises

H) reacting a compound of formula (7)




wherein R is as defined above, with a compound of formula (2) as defined above, to obtain a compound of formula (8) as defined above; and further submitting the resulting compound of formula (8) first to step (K) and then to step (N) as disclosed above.

Process step (H) may preferably be effected in the presence of sodium ethylate, e.g. as disclosed for step (d) above.


Compound of formula (7) used in process step (H) may be prepared by

c) iodinating a compound of formula (19)



wherein R is as defined above; or

- l) reducing a compound of formula (7')




wherein R' is as defined above.


Process step (c) may conveniently be performed by reaction with an iodinating agent, e.g. an alkali metal iodide such as NaI or KI. Process step (l) may be carried out under acidic conditions, e.g. trifluoroacetic acid. Suitable reducing agents include e.g. triethylsilane.

Compound of formula (19), used as starting product in the preparation of 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, is novel and also forms part of the present invention as well as its preparation. Compound of formula (19) may be obtained by

- C) hydrogenating a compound of formula (10)



wherein R' is as defined above, to obtain a compound of formula (5)



wherein R is as defined above; and

- C') brominating the resulting compound of formula (5).

Process step (C) may be performed as disclosed e.g. for (j). Process step (C') may be carried out in accordance with methods known in the art, e.g. using a brominating agent, e.g. HBr, in the presence of a phase transfer-catalyst, e.g. an ammonium salt such as tricaprylmethylammonium halogenide, and further purification using e.g. aluminium oxide.

Compound of formula (19) may also be prepared by

- b) reducing a compound of formula (18) as defined above.

Process step (b) may be carried out as disclosed for step (l) above.

Compound of formula (19) is useful to prepare the intermediary compound of formula (8) used in process step (K)

- e) directly by reaction with a compound of formula (2) as defined above, preferably in the presence of a base, e.g. an alkali metal hydride such as NaH, or an alkali metal alkoxide, e.g. C_2H_5ONa or $t-C_4H_9OK$.

Process step e) may be carried out in accordance with known methods or e.g. as disclosed in Example 20.

According to the invention, Compound A is preferably prepared either starting from a compound of formula (19) and following above disclosed route: step (e) \Rightarrow step (K) \Rightarrow step (N); or starting from a compound of formula (7) and following above disclosed route: step (H) \Rightarrow step (K) \Rightarrow step (N); or starting from a compound of formula (20) or (20') and following above disclosed route: step (f) \Rightarrow step (K) \Rightarrow step (N).

The present invention also comprises each of the following steps disclosed above, respectively: step (b), step (c), step (C'), step (e), step (d), step (g'), step (g), step (x), step (f).

Examples of salts include salts with inorganic acids, such as hydrochloride, hydrobromide, and sulfate, salts with organic acids, such as acetate, lactate, succinate or tartarate, fumarate, maleate, benzoate, citrate, malate, methanesulfonate and benzenesulfonate salts. Preferred salt is the hydrochloride salt. 2-amino-2-[2-(4- C_{2-20} alkylphenyl)ethyl]-propane-1,3-diols, preferably Compound A, may be converted into the hydrochloride salt form in accordance with known methods, e.g. by addition of HCl to the last reaction step such as in step (j), or prior to recrystallization such as in step (N).

2-Amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diols and/or the intermediate compounds may be purified and/or separated by a conventional manner such as recrystallization, column chromatography, distillation, centrifugal separation, washing or drying.

The present Examples are illustrative of the invention.

Example 1: Production of 2-acetamido-4-acetoxy-2-acetoxymethyl-4-(4-octylphenyl)butyl acetate (21) [Step (g)]

To a solution of diethyl acetamido-2-(4-octylphenyl)-2-oxoethylmalonate (20) (2.00 g) in methanol (8 ml) is added sodium borohydride (858 mg) and the mixture is stood at room temperature for 1.5 hr. The suspension is diluted with ethyl acetate and washed successively with 1N HCl, saturated aqueous NaHCO₃ solution and saturated brine. The ethyl acetate layer is dried over anhydrous MgSO₄ and concentrated. To the residue is added pyridine (4 ml)-acetic anhydride (8.8 ml), and the mixture is stood at room temperature overnight. To the reaction mixture is added ice water and the mixture is extracted with ethyl acetate and washed successively with 1N HCl, saturated aqueous NaHCO₃ solution and saturated brine. The ethyl acetate layer is dried over anhydrous MgSO₄ and concentrated. The residue is purified by silica gel column chromatography using hexane-ethyl acetate as eluent to give the title compound (21) as a colorless oil.

TLC Rf: 0.3 (hexane/ethyl acetate= 1/2, silica gel 60F₂₅₄ plate)

IR (CCl₄) 3390, 2930, 2860, 1750, 1690, 1510, 1370, 1230, 1040 cm⁻¹.

¹H-NMR (500 MHz, CDCl₃) δ: 7.22 (2H, d, J=8.1Hz, C₆H₂), 7.14 (2H, d, J=8.1 Hz, C₆H₂), 5.91 (1H, brs, NH), 5.84 (1H, dd, J=4.2 and 8.5Hz, CH), 4.37 (1H, d, J= 11.2Hz, OCHa), 4.34 (1H, d, J= 11.2Hz, OCHb), 4.34 (2H, s, OCH₂), 2.56 (2H, t, J=7.8Hz, PhCH₂), 2.54 (1H, dd, J=4.2 and 15.1Hz, CHCHa), 2.36 (1 H, dd, J=8.5 and 11.5Hz, CHCHb), 2.10, 2.07 and 1.99 (each 3H, s, OAc), 1.90 (3H, s, NAc), 1.57 (2H, qui, J=7.8Hz, CH₂), 1.32-1.23 (10H, m, CH₂ x 5), 0.87 (3H, t, J=7.1 Hz, CH₃)

FAB-MS m/z: 432 (M-AcOH+H)⁺

Example 2: Production of 2-amino-2-(2-hydroxy-2-(4-octylphenyl)ethyl)propane-1,3-diol (22) [Step (f)]

- 11 -

2-Acetamido-4-acetoxy-2-acetoxymethyl-4-(4-octylphenyl)butyl acetate (21) (700 mg) is refluxed under heating for 5 hr in methanol (3.6 ml)/1N NaOH (7.2 ml). The reaction mixture is diluted with water and subjected to XAD-II column using water and methanol as eluents. The methanol eluate is concentrated to give the title compound (22) as a pale-yellow solid. TLC Rf:0.5 (chloroform/methanol/acetic acid/water=70/20/6/4, silica gel 60F₂₅₄ plate) IR (KBr) 3340, 2930, 2850, 1470, 1040 cm⁻¹.

¹H-NMR (500 MHz, DMSO-d₆) δ: 7.20 (2H, d, J=8.1Hz, C₆H₂), 7.10 (2H, d, J=8.1Hz, C₆H₂), 6.66 (1H, brs, CHOH), 4.83 (1H, dd, J=3.0 and 10.0Hz, CH), 4.72 and 4.61 (each 1H, brs, OH), 3.42 (2H, s, OCH₂), 3.25 (1H, d, J=10.5Hz, OCHa), 3.21 (1H, d, J= 10.3Hz, OCHb), 2.52 (2H, t, J=7.6Hz, PhCH₂), 1.71 (2H, brs, NH₂), 1.55-1.51 (2H, m, CH₂), 1.51 (1H, dd, J= 3.2 and 14.4 Hz, CHCHa), 1.47 (1H, dd, J= 10.1 and 14.4Hz, CHCHb), 1.27-1.23 (10H, m, CH₂ X 5), 0.85 (3H, t, J=7.1Hz, CH₃)

FAB-MS m/z: 324 (M+H)⁺

Example 3: Production of 2-amino-2-[2-(4-octylphenyl)ethyl]propane-1,3-diol hydrochloride (Compound A in hydrochloride salt form) [Step (I)]

5% Pd/carbon (13 mg) is suspended in ethanol (0.5 ml) and a solution of 2-amino-2-[2-hydroxy-2-(4-octylphenyl)ethyl]propane-1,3-diol (22) (107 mg) in ethanol (4 ml)-1N HCl in ethanol (0.55 ml) is added. The mixture is stirred at room temperature for 10 days under 5 kg/cm² hydrogen pressurization. The reaction mixture is filtered and the filtrate is concentrated to Compound A (hydrochloride) as colorless crystals.

Decomposition: 260°C

TLC Rf : 0.55 (chloroform/methanol/ acetic acid/water=70/20/6/4, silica gel 60F₂₅₄ plate)

IR (KBr) 3400 (sh), 3250, 3050 (sh), 2910, 2850, 1580, 1520, 1470, 1060 cm⁻¹.

UVλ_{max}(H₂O) nm (ε): 210.7(4709), 264(392.4), 272(341.1)

¹H-NMR (500 MHz, DMSO-d₆) δ: 7.91 (3H, br s; NH₃⁺), 7.09 (2H, d, J=8.5Hz, C₆H₂), 7.07 (2H, d, J=8.5Hz, C₆H₂), 5.38 (2H, br s, OH x 2), 3.51 (4H, s, CH₂OH x 2), 2.56 (2H, m, Ph-CH₂), 2.49 (2H, m, Ph-CH₂), 1.77 (2H, m, CH₂), 1.51 (2H, m, CH₂), 1.25 (10H, m, CH₂ x 5), 0.83 (3H, t, J=7.5Hz, CH₃)

EIMS m/z: 276 (M- CH₂OH)⁺, 117, 105

Example 4: Production of 2-acetamido-2-acetoxymethyl-4-(4-octylphenyl)butyl acetate (9)
[Step (h)]

5% Pd/carbon (20 mg) is suspended in ethyl acetate (0.5 ml) and a solution of 2-acetamido-4-acetoxy-2-acetoxymethyl-4-(4-octylphenyl)butyl acetate (21) (200 mg) in ethyl acetate (5 ml) and acetic acid (0.2 ml) are added. The mixture is stirred at room temperature for 6 days under 5 kg/cm² hydrogen pressurization. The reaction mixture is filtered and the filtrate is washed successively with saturated aqueous NaHCO₃ solution and saturated brine. The ethyl acetate layer is dried over anhydrous MgSO₄ and concentrated to give the title compound (9) as colorless crystals. m.p. 111.8°C

TLC Rf:0.4 (hexane/ethyl acetate = 1/2, silica gel 60F₂₅₄ plate)

IR (KBr) 3330, 2910, 2850, 1740, 1650, 1550, 1470, 1390, 1260, 1240, 1050cm⁻¹.

UVλ_{max}(MeOH) nm (ε): 217.6(4772), 259.0(305.7), 264.5(394.6), 272.8(368.6)

¹H-NMR (270 MHz, DMSO-d₆) δ: 7.63 (1H, br s, NH), 7.07 (4H, s, C₆-H₄), 4.28 (2H, d, J= 10.6Hz, CHaOAc x 2), 4.18 (2H, d, J= 10.6Hz, CHbOAc x 2), ca. 2.5 (4H, m, Ph-CH₂ x 2), 2.02 (6H, s, OCOCH₃, x 2), 1.94 (2H, m, CH₂), 1.85 (3H, s, NCOCH₃), 1.52 (2H, m, CH₂), 1.24 (10H, m, CH₂ x 5), 0.85 (3H, t, J=7.2Hz, CH₃)

EIMS m/z: 433 (M)⁺, 373, 360, 300, 216, 157, 117, 105, 97

Example 5: Production of 2-amino-2-[2-(4-octylphenyl)ethyl]propane-1,3-diol hydrochloride (Compound A in hydrochloride salt form) [Step (N)]

2-Acetamido-2-acetoxymethyl-4-(4-octylphenyl)butyl acetate (9) (7.37 kg) is refluxed under heating in methanol (51 L)-2N aqueous NaOH solution (34 L) for 6 hr. After cooling, the resulting precipitate is collected by filtration and washed thoroughly with water. The precipitate is dissolved in conc. HCl/ethanol (1:11) (20 L) and concentrated. The residue is recrystallized from ethyl acetate/ethanol to give Compound A (hydrochloride) as colorless crystals. The resulting Compound A has the same physico-chemical characteristics as indicated in Example 3.

Example 6: Production of 2-acetamido-2-acetoxymethyl-4-(4-octylphenyl)butyl acetate (9)
[Step (x)]

To a solution of diethyl acetamido-2-(4-octylphenyl)-2-oxoethylmalonate (20) (200 mg) in methanol (0.8 ml) is added sodium borohydride (85 mg) and the mixture is stood at room temperature for 2.5 hr. The reaction mixture is diluted with ethyl acetate and water, 1N HCl (2.24 ml) and acetic acid (0.26 ml) are added, followed by partitioning. The ethyl acetate layer is washed successively with saturated aqueous NaHCO_3 solution and saturated brine. The ethyl acetate layer is dried over anhydrous MgSO_4 and concentrated. The residue is stirred in ethanol (2 ml) in the presence of 5% Pd/ Carbon (20 mg) and acetic acid (0.2 ml) for 8 days under 5 kg/cm² hydrogen pressurization. The reaction mixture is filtered through a membrane filter and the filtrate is concentrated.

The residue is dissolved in acetic anhydride (2.8 ml)-pyridine (1 ml), and the mixture is stood at room temperature overnight. The reaction mixture is poured into ice water and the precipitate is collected by filtration. The precipitate is recrystallized from hexane-ethyl acetate to give the title compound (9) as colorless crystals. The above-mentioned recrystallization filtrate is concentrated and purified by preparative TLC to give further the title compound (9) as colorless crystals. The resulting Compound (9) has the same physico-chemical characteristics as indicated in Example 4.

Alternatively compound (20) may be hydrogenated in ethanol in the presence of 5% Pd/C and then treated with NaBH_4 and then with acetic anhydride-pyridine as disclosed above.

Example 7: Production of diethyl acetamido-2-(4-octylphenyl)ethylmalonate (8) [Step (f)]

Diethyl acetamido-2-(4-octylphenyl)-2-oxoethylmalonate (20) (200 mg) is stirred at room temperature for 6 days in ethanol (2 ml) in the presence of 5% Pd/carbon (20 mg) under 5 kg/cm² hydrogen pressurization. The reaction mixture is filtered and the filtrate is concentrated. The residue is dissolved in ethyl acetate and washed successively with saturated aqueous NaHCO_3 solution and saturated brine. The ethyl acetate solution is dried over anhydrous MgSO_4 and concentrated to give the title compound (8). m.p. 61°C

Alternatively diethyl acetamido-2-(4-octylphenyl)ethylmalonate (8) is prepared by hydrogenation of diethyl acetamido-2-(4-octan-1'-ylphenyl)-ethylmalonate in the presence of 5% Pd/carbon as disclosed above.

Compound (8) has the same physico-chemical characteristics as indicated above.

TLC Rf : 0.6 (hexane/ethyl acetate = 1/1, silica gel 60F₂₅₄ plate).

IR (KBr) 3300, 2920, 2850, 1750, 1650, 1520, 1220, 1200 cm⁻¹.

UVλ_{max}(MeOH) nm (ε): 219.1(5017), 259.2(303.5), 264.5(392.4), 272.7(357.7).

¹H-NMR (270 MHz, DMSO-d₆) δ: 8.32 (1H, br s, NH), 7.08 (2H, d, J=7.9Hz, C₆-H₂), 7.02 (2H, d, J=7.9 Hz, C₆-H₂), 4.13 (4H, q, J=7.3 Hz, OCH₂CH₃ X 2), 2.52 (4H, m, Ph-CH₂ X 2), 2.37 (2H, m, CH₂), 1.94 (3H, s, COCH₃), 1.52 (2H, m, CH₂), 1.24 (10H, m, CH₂ X 5), 1.15 (6H, t, J=7.3Hz OCH₂CH₃ X 2), 0.85 (3H, t, J=6.6Hz, CH₃)

EIMS m/z: 388 (M-OCH₂CH₃)⁺, 318, 301, 244, 217, 171, 143.

Example 8: Production of 2-acetamido-2-acetoxymethyl-4-(4-octylphenyl)butyl acetate (9)
[Step (K)]

To a solution of diethyl acetamido-2-(4-octylphenyl)ethylmalonate (8) (1.37 kg) in methanol (5.5 L) is added sodium borohydride (720 g), and the mixture is stirred at room temperature for 2 hr. The suspension is diluted with ethyl acetate, and washed successively with 4N HCl, saturated aqueous NaHCO₃ solution and saturated brine. The ethyl acetate layer is dried over anhydrous magnesium sulfate and concentrated. To the residue are added pyridine (3 L) and acetic anhydride (12 L), and the mixture is stirred at room temperature overnight. The reaction mixture is poured into ice water and the precipitated crystals are collected by filtration. The precipitate is recrystallized from hexane/ethyl acetate (2/1) to give the title compound (9) as colorless crystals. The resulting Compound (9) has the same physico-chemical characteristics as indicated in Example 4.

Alternatively, the reduction of Compound (8) may be carried out using NaBH₄ and CaCl₂, e.g. in aqueous ethanol.

Example 9: Production of 2-acetamido-2-[2-hydroxy-2-(4-octylphenyl)ethyl]propane-1,3-diol (21') [Step (g')]

Diethyl 2-acetamido-2-[2-(4-octylphenyl)-2-oxoethyl]malonate (20) (4 g) is dissolved in methanol (150 ml) and sodium borohydride (tablet, 1.64 g) is added. The mixture is stirred for 1 hr under heating at 50°C. The solvent is evaporated and the residue is extracted with

ethyl acetate. The extract is washed with brine and dried over sodium sulfate. The solvent is evaporated to give the title compound (21') as colorless crystals. m. p. 118-119°C
IR (KBr): 3228, 2927, 1640, 1566, 1071 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3): 0.86 (3H, t, $J=6\text{Hz}$), 1.18-1.33 (10H, m), 1.48-1.64 (2H, m), 1.57 (2H, bs), 1.80 (1H, dd, $J=8\text{Hz}$, $J=10\text{Hz}$), 2.02 (3H, s), 2.34 (1H, d, $J=10\text{Hz}$), 2.56 (2H, t, $J=6\text{Hz}$), 3.39-3.47 (1H, m), 3.54-3.62 (1H, m), 3.68-3.80 (2H, m), 4.11 (1H, br. s), 4.85 (1H, d, $J=7\text{Hz}$), 6.96 (1H, br. s), 7.14 (2H, d, $J=8\text{Hz}$), 7.23 (2H, $J=8\text{Hz}$)
MS (EI) m/z : 365 (M^+)

Example 10: Production of 2-amino-2-[2-hydroxy-2-(4-octylphenyl)ethyl]propane-1,3-diol (22) [Step (I')]

2-Acetamido-4-(4-octylphenyl)-2-hydroxymethylbutane-1,4-diol (21') (0.73 g) is dissolved in methanol (10 ml). Thereto is added a solution of LiOH (0.86 g) in water (10 ml), and the mixture is refluxed under heating for 2 hr. The reaction mixture is cooled and extracted with ethyl acetate. The organic layer is washed with brine and dried over sodium sulfate. The solvent is distilled away and the mixture is purified by silica gel column chromatography to give the title compound (22) as a colorless and amorphous powder. The resulting Compound (22) has the same physico-chemical characteristics as indicated in Example 2.

Example 11: Production of 2-bromo-4'-octylacetophenone (18) [Step (a)]

To a suspension of aluminum chloride (3.88 g) and bromoacetyl chloride (4.56 g) in dichloromethane (5 ml) is added dropwise over 1 hr a solution of octyl-benzene (16) (5.0 g) in dichloromethane (7 ml) under ice-cooling, and the mixture is stirred under ice-cooling for 1 hr. The reaction mixture is poured into ice water and extracted with ethyl acetate. The organic layer is washed successively with HCl, saturated aqueous NaHCO_3 solution and saturated brine, dried over anhydrous MgSO_4 and concentrated to give the title compound (18) as a colorless oil.

TLC Rf: 0.4 (hexane/ethyl acetate=20 : 1, silica gel 60F₂₅₄ plate).
IR (CCl_4) 2930, 2860, 1680, 1610, 1430, 1280, 1180, 1010 cm^{-1} .

- 16 -

¹H-NMR (500 MHz, CDCl₃) δ: 7.90 (2H, d, J=8.5Hz, C₆H₂), 7.29 (2H, d, J=8.5Hz, C₆H₂), 4.44 (2H, s, BrCH₂), 2.67 (2H, t, J=7.6Hz, PhCH₂), 1.63 (2H, qui, J=7.1Hz, CH₂), 1.32-1.26 (10H, m, CH₂x5), 0.88 (3H, t, J=7.1Hz, CH₃)

¹³C-NMR (500 MHz, CDCl₃) δ: 190.93, 149.95, 131.60, 129.05, 128.89, 36.06, 31.82, 31.01, 30.97, 29.38, 29.24, 29.19, 22.63, 14.08

EIMS m/z: 312 and 310 (M)⁺, 217, 91

Example 12: Production of 2-bromo-4'-octylacetophenone (18)

4'-Octylacetophenone (18') [Step (a'')]

Aluminum chloride (22.6 g) is suspended in dichloroethane (500 ml) and octylbenzene (16) (21.5 g) is added at room temperature. To the mixture is added acetyl chloride (8.87 g) under ice-cooling. The mixture is stirred at room temperature overnight. The reaction mixture is poured into ice water and extracted with ether. The organic layer is washed with brine and dried over sodium sulfate. The solvent is evaporated to give 4'-octylacetophenone (18') as a colorless oil.

2-Bromo-4'-octylacetophenone (18) [Step (a''')]

To a solution of 4'-octylacetophenone (18') (23.9 g) in acetic acid (100 ml) is added hydrogen bromide-acetic acid solution (25%, 34 ml) under ice-cooling, which is followed by dropwise addition of bromine (5.3 ml). The mixture is heated to 30°C and stirred for 3 hr. The solvent is distilled away and the mixture is purified by silica gel column chromatography to give 2-bromo-4'-octylacetophenone as colorless crystals.

m. p. 33-34°C.

The resulting Compound (18) has the same physico-chemical characteristics as indicated in Example 11.

Example 13: Production of diethyl acetamido-2-(4-octylphenyl)-2-oxoethylmalonate (20) [Step (d)]

A solution of 2-bromo-4'-octylacetophenone (18) (5.0 g), diethyl acetamidomalonate (3.85 g) and sodium ethylate (1.20 g) in anhydrous ethanol (50 ml) is refluxed under heating under nitrogen atmosphere for 6 hr. The reaction mixture is diluted with hexane/ethyl

- 17 -

acetate, washed with water, dried over anhydrous MgSO_4 and concentrated. The residue is purified by silica gel column chromatography using hexane/ethyl acetate as eluent to give the title compound (20) as a pale-yellow oil.

TLC Rf: 0.5 (hexane/ethyl acetate=1/1, silica gel 60F₂₅₄ plate)

IR (CCl_4): 3420, 2930, 2860, 1750, 1690, 1610, 1490, 1350, 1230, 1200 cm^{-1} .

$^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 7.87 (2H, d, $J=8.3\text{Hz}$, C_6H_2), 7.25 (2H, d, $J=8.3\text{Hz}$, C_6H_2), 7.11 (1 H, s, NH), 4.26 and 4.40 (each 2H, q, $J=7.1\text{ Hz}$ OCH_2), 4.23 (2H, s, COCH_2), 2.65 (2H, t, $J=7.8\text{Hz}$, PhCH_2), 1.96 (3H, s, Ac), 1.61 (2H, qui, $J=7.3\text{Hz}$, CH_2), 1.30-1.22 (10H, m, $\text{CH}_2 \times 5$), 1.23 (6H, t, $J=7.1\text{Hz}$, $\text{CH}_3 \times 2$), 0.87 (3H, t, $J=7.1\text{ Hz}$, CH_3)

FAB-MS m/z : 448 ($\text{M}+\text{H}$)⁺

Example 14: Production of 2-(4-octylphenyl)ethyl bromide (19) [Step (b)]

To a solution of 2-bromo-4'-octylacetophenone (18) (1.0 g) in trifluoroacetic acid (2.5 ml) is added triethylsilane (1.12 ml) under ice-cooling, and the mixture is stirred at room temperature for 2 hr. The reaction mixture is poured into ice water and hexane and saturated aqueous NaHCO_3 solution are added, which is followed by partitioning. The hexane layer is washed with saturated brine, dried over anhydrous MgSO_4 and concentrated. The residue is purified by silica gel column chromatography using hexane as eluent to give the title compound (19) as a colorless oil.

TLC Rf:0.5 (hexane, silica gel 60F₂₅₄ plate)

IR (CCl_4) 2930, 2860, 1510, 1470 cm^{-1} .

$^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 7.15 (2H, d, $J=8.3\text{Hz}$, C_6H_2), 7.12 (2H, d, $J=8.3\text{Hz}$, C_6H_2), 3.56 (2H, t, $J=7.8\text{Hz}$, BrCH_2), 3.14 (2H, t, $J=7.8\text{Hz}$, PhCH_2), 2.59 (2H, t, $J=7.8\text{Hz}$, PhCH_2), 1.61 (2H, qui, $J=7.6\text{Hz}$, CH_2), 1.35-1.27 (10H, m, $\text{CH}_2 \times 5$), 0.89 (3H, t, $J=7.3\text{Hz}$, CH_3)

EIMS m/z : 298 and 296 (M)⁺, 217, 197 and 199, 117, 105, 91

Example 15: Production of 2-(4-octylphenyl)ethyl iodide (7) [Step (c)]

To a solution of 2-(4-octylphenyl)ethyl bromide (19) (99 mg) in 2-butanone (5 ml) is added NaI (126mg) and the mixture is refluxed under heating for 3 hr. The reaction mixture is diluted with hexane and washed successively with saturated aqueous sodium sulfite

- 18 -

solution and saturated brine. The hexane layer is dried over anhydrous Mg SO_4 and concentrated to yield the title compound (7) as a colorless oil.

TLC Rf:0.6 (hexane, silica gel 60F₂₅₄ plate)

IR (CCl_4) 2930, 2860, 1510, 1470, 1170 cm^{-1} .

$^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 7.13 (2H, d, $J=8.1\text{Hz}$, C_6H_2), 7.10 (2H, d, $J=8.1\text{Hz}$, C_6H_2), 3.33 (2H, t, $J=7.6\text{Hz}$, ICH_2) 3.15 (2H, t, $J=7.8\text{Hz}$, PhCH_2), 2.57 (2H, t, $J=7.8\text{Hz}$, PhCH_2), 1.60 (2H, qui, $J=7.8\text{Hz}$, CH_2), 1.34-1.26 (10H, m, $\text{CH}_2 \times 5$), 0.88 (3H, t, $J=6.6\text{Hz}$, CH_3)

EIMS m/z : 344 (M^+), 245, 217, 119, 117, 91, 57, 43

Example 16: Production of diethyl acetamido-2-(4-octylphenyl)ethylmalonate (8) [Step (H)]

To a solution of diethyl acetamidomalonate (277 g) and sodium ethylate (86.6 g) in anhydrous ethanol (850 ml) is added a solution of 2-(4-octylphenyl)ethyl iodide (7) (146 g) in anhydrous tetrahydrofuran (533 ml), and the mixture is refluxed under heating for 6 hr. The reaction mixture is concentrated and partitioned between water and hexane, and the hexane layer is washed three times with water. The obtained hexane layer is dried over anhydrous magnesium sulfate and concentrated. The residue is recrystallized from hexane to give the title compound (8) as colorless crystals. The resulting Compound (8) has the same physico-chemical characteristics as indicated in Example 7.

Example 17:: Production of 2-(4-octylphenyl)ethyl iodide (7) [Step (I)]

To a solution of 4'-(2-iodoethyl)octanophenone (200 g) in trifluoroacetic acid (319 ml) is added triethylsilane (195.7 ml) under ice-cooling, and the mixture is stirred at room temperature for 2 hr. The reaction mixture is concentrated, and the residue is distilled away under reduced pressure to give the title compound (7) as a pale-red oil. The resulting Compound (7) has the same physico-chemical characteristics as indicated in Example 15.

Example 18: Production of 2-(4-octylphenyl)ethanol (5) [Step (C)]

5% Pd/carbon (0.6 g) is added to a solution of 2-(4-octanoylphenyl)ethanol (10) (30 g) in abs. ethanol (150 ml). The mixture is stirred at room temperature for 2.5 hrs under 0.1 bar

- 19 -

hydrogen. The reaction mixture is filtered and the filtrate is concentrated to afford compound (5) as a clear pale yellow oil.

IR (NaCl): 3300, 2926, 2854, 1513, 1466, 1046 cm^{-1} .

$^1\text{H-NMR}$ (200 MHz, CDCl_3) δ : 7.12 (4H, s, C_6H_4); 3.87 (2H, t, $J=7.7\text{Hz}$, CH_2O); 2.85 (2H, t, $J=7.7\text{Hz}$, Ph-CH_2); 2.59 (2H, t, $J=7.8\text{Hz}$, Ph-CH_2); 1.61 (2H, q, $J=7.8\text{Hz}$, CH_2); 1.40-1.15 (10H, m, $\text{CH}_2 \times 5$); 0.89 (3H, t, 7.3Hz, CH_3)

EIMS m/z : 234 (M^+), 203, 135, 105

Example 19: Production of 2-(4-octylphenyl)ethyl bromide (19) [Step (C')]

2-(4-octylphenyl)ethanol (5) (46.9 g) is added within 10 min to an aqueous solution containing 3.05 g Na_2SO_3 , 252.9 g 48% HBr and 3.2 g tricaprylmethylammonium chloride. The reaction mixture is heated to 100°C for 18 to 22 hours with stirring. The resulting 2-phase mixture is cooled to room temperature, diluted with toluene (150 ml) and the aqueous phase is then removed. The organic phase is washed twice with a volume of 100 ml of a 9% NaHCO_3 solution. The organic phase is treated with basic aluminium oxide, then filtered. The filter residue is washed with toluene (50 ml) and evaporated to yield compound (19) as a clear pale yellow oil. The resulting Compound (19) has the same physico-chemical characteristics as indicated in Example 14.


Example 20: Production of diethyl acetamido-2-(4-octylphenyl)ethylmalonate (8) [Step (e)]

60% NaH in oil (9.65 g) is added portionwise to a solution of diethyl acetamidomalonate (58.5 g) in dimethylformamide (240 ml) at such a rate to keep the temperature below 10°C . The resulting mixture is stirred for 90 min., the temperature being allowed to increase to room temperature. To this mixture is added dropwise within 15 min. a solution of 2-(4-octylphenyl)ethyl bromide (19) (40 g) in dimethylformamide (40 ml). The reaction mixture is heated to ca. 80°C within ca. 25 min. and further stirred for ca. 3-5 hrs. After cooling to room temperature, water (280 ml) is added and the resulting mixture is further stirred until a beige precipitate is formed. The mixture is adjusted to pH 7 with 10% H_2SO_4 . After cooling to ca. 2°C , the residue is filtered, washed with 200 ml water and then recrystallized in heptane to yield compound (8) as colorless crystals. The resulting compound (8) exhibits the same physico-chemical characteristics as indicated in Example 7.


- 20 -

CLAIMS


1. A process for preparing a 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises
- g) reducing a compound of formula (20)



wherein R is C₂₋₂₀alkyl, each of R₁ and R₂, independently, is C₁₋₄alkyl and R₃ is a protecting group,
in the presence of (R₄-CO)₂O wherein R₄ is C₁₋₄alkyl, to obtain a compound of formula (21)



- wherein R, R₃ and R₄ are as defined above; and
- i) treating the resulting compound of formula (21) with a basic compound to obtain a compound of formula (22)




wherein R is as defined above; and

- 21 -

j) hydrogenating the resulting compound (22) and, where required, converting the resulting 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, into the desired salt form.

2. A process according to claim 1, comprising

h) hydrogenating a compound of formula (21) to obtain a compound of formula (9)



wherein R, R₃ and R₄ are as defined in claim 1; and

N) treating the resulting compound of formula (9) with a basic compound and, where required, converting the resulting 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, into the desired salt form.


3. A process for preparing a 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises

x) reducing a compound of formula (20) as defined in claim 1, followed by a treatment with (R₄CO)₂O, to obtain a compound of formula (9) as defined in claim 2; and

N) treating the resulting compound of formula (9) with a basic compound and, where required, converting the resulting 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, into the desired salt form.


4. A process for preparing a 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises

f) hydrogenating a compound of formula (20) as defined in claim 1 or a compound of formula (20')



- 22 -

wherein R_1 , R_2 and R_3 are as defined in claim 1, and R' is C_{1-19} -CO- to obtain a compound of formula (8)




wherein R and R_1 to R_3 are as defined in claim 1; and

- K) reducing the resulting compound of formula (8) followed by a treatment with $(R_4CO)_2O$ to obtain a compound of formula (9) as defined in claim 2; and
- N) treating the resulting compound of formula (9) with a basic compound and, where required, converting the resulting 2-amino-2-[2-(4- C_{2-20} alkylphenyl)ethyl]propane-1,3-diol, into the desired salt form.

5. A process for preparing a 2-amino-2-[2-(4- C_{2-20} alkylphenyl)ethyl]propane-1,3-diol, in free form or in salt form, which process comprises


- g') reducing a compound of formula (20) as defined in claim 1 under conditions to obtain a compound of formula (21')



wherein R and R_3 are as defined in claim 1; and

- y) hydrogenating the resulting compound of formula (21') to obtain a compound of formula (21'')

- 23 -



wherein R and R₃ are as defined in claim 1; and

- z) treating the resulting compound of formula (21') with a basic compound and, where required, converting the resulting 2-amino-2-[2-(4-C₂₋₂₀alkylphenyl)ethyl]propane-1,3-diol, into the desired salt form.


6. A process according to claim 5, comprising
 i') reacting a compound of formula (21') with a basic compound to obtain a compound of formula (22) as defined in claim 1

which compound of formula (22) is then further treated as defined in claim 1.

7. A process for preparing a compound of formula (20) as defined in claim 1, comprising
 a) reacting a compound of formula (16)




wherein R is C₂₋₂₀ alkyl, with 2-bromoacetyl chloride to obtain a compound of formula (18)



wherein R is C₂₋₂₀alkyl; and


- d) reacting the resulting compound of formula (18) with a compound of formula (2)

- 24 -




wherein R_1 to R_3 are as defined in claim 1.

8. A process according to claim 7, comprising
 a') reacting a compound of formula (16) as defined in claim 7 above, with acetyl chloride to obtain a compound of formula (18')




wherein R is C_{2-20} alkyl,
 which compound of formula (18') is brominated to give a compound of formula (18).

9. A process for preparing a 2-(4- C_{2-20} alkylphenyl)ethyl bromide comprising
 b) reducing a compound of formula (18)



wherein R is C_{2-20} alkyl.


10. A process for the preparation of a compound of formula (8)



wherein R is C_{2-20} alkyl, each of R_1 and R_2 , independently, is C_{1-4} alkyl and R_3 is a protecting group, comprising


- b) reducing a compound of formula (18) as defined in claim 9 to obtain a compound of formula (19)

- 25 -




wherein R is C₂₋₂₀ alkyl, and

- c) iodinating the resulting compound of formula (19), to obtain a compound of formula (7)




wherein R is C₂₋₂₀ alkyl, and

- H) reacting the resulting compound of formula (7) with a compound of formula (2)



wherein R₁ to R₃ are as defined in claim 1.


11. A process according to any of claims 1 to 6 for the preparation of 2-amino-2-[2-(4-octylphenyl)ethyl]propane-1,3-diol, in free form or in salt form.
12. A compound of formula (20)



wherein R is C₂₋₂₀ alkyl, each of R₁ and R₂, independently, is C₁₋₄ alkyl and R₃ is a protecting group.

13. A compound according to claim 14 which is diethyl acetamido-2-(4-octylphenyl)-2-oxoethylmalonate.

14. A compound of formula (19)




wherein R is C₂₋₂₀ alkyl.


15. A compound according to claim 14 which is 2-(4-octylphenyl)ethyl bromide.

16. A process for preparing a compound of formula (19) as defined in claim 14, comprising

- C) hydrogenating a compound of formula (10)




wherein R' is C₁₋₁₉alkyl-C=O, to obtain a compound of formula (5)



wherein R is C₂₋₂₀alkyl; and

- C') brominating the resulting compound of formula (5).

17. A process according to claim 4 comprising the step of
e) reacting a compound of formula (19) as defined in claim 14 with a compound of formula (2)



wherein R₁ to R₃ are as defined in claim 4, to obtain a compound of formula (8) as defined in claim 4.

18. A process according to claim 17 for the preparation of 2-amino-2-[2-(4-octylphenyl)ethyl]propane-1,3-diol in free form or in salt form, comprising the step of reacting 2-(4-octylphenyl)ethyl bromide with diethyl acetamidomalonate.
19. A process according to any one of claims 1 to 10, 16 and 17, wherein each of C₂-alkyl and R is octyl.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/08577

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07C215/28 C07C213/08 C07C22/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DD 269 856 A (UNIV HALLE WITTENBERG) 12 July 1989 (1989-07-12) page 3; figure A	14
X	--- "BEILSTEINS HANDBUCH DER ORGANISCHEN CHEMIE, Vol. V, 4. Edition, Suppl. 3, Page 970" 1964, SPRINGER VERLAG, BERLIN. GÖTTINGEN. HEIDELBERG. NEW YORK XP002127759 paragraph [0002] --- -/-	14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

18 January 2000

Date of mailing of the international search report

27. 4. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

O'Sullivan, P

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/08577

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 100, no. 22, 28 May 1984 (1984-05-28) Columbus, Ohio, US; abstract no. 192463f, KESZLER, B.; CHANG, V. S. C.; KENNEDY, J. P.: "Styryl-telechelic polyisobutylenes. I. Synthesis of linear and tri-arm star styryl-telechelic polyisobutylenes." page 12; column 1; XP002127760 abstract & J. MACROMOL. SCI., CHEM., vol. A21, no. 3, 1984, pages 315-326, ---	14
X	BERNSTEIN, P. R.; SNYDER, D. W.; ADAMS, E. J.; KRELL, R. D.; VACEK, E. P.; WILLARD, A. K.: "Chemically Stable Homocinnamyl Analogues of the Leukotrienes: Synthesis and Preliminary Biological Evaluation." J. MED. CHEM., vol. 29, 1986, pages 2477-2483, XP000867024 page 2478; figure 1 ---	14
A	EP 0 627 406 A (YOSHITOMI PHARMACEUTICAL) 7 December 1994 (1994-12-07) cited in the application page 166; example 28 page 8, line 10 page 168; example 29 page 7, line 40 -----	1-8, 10-15, 17,18

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/ 08577

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-8, 10-15, 17, 18

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/EP 99/08577

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-8, 10-15, 17, 18

Process for the production of 2-amino-2-(2-(4-C₂-20-alkyl-phenyl)ethyl)propane-1,3 diols having as intermediate a derivative of compound 20

2. Claim : 9

Process for the preparation of compound (19)
2-(4-alkylphenyl)ethyl bromide from compound (18)

3. Claim : 16

Process for the preparation of compound (19)
2-(4-alkylphenyl)ethyl bromide from compound (10)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internatic Application No
PCT/EP 99/08577

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DD 269856	A	12-07-1989	NONE	
EP 0627406	A	07-12-1994	DE 69321823 D	03-12-1998
			DE 69321823 T	02-06-1999
			US 5604229 A	18-02-1997
			AT 172711 T	15-11-1998
			CA 2126337 A	28-04-1994
			ES- 2126658 T	01-04-1999
			WO 9408943 A	28-04-1994
			JP 2579602 B	05-02-1997
			KR 155015 B	01-12-1998
			US 5719176 A	17-02-1998
			US 5952316 A	14-09-1999